

THE NORWEGIAN UNIVERSITY OF SCIENCE AND TECHNOLOGY
DEPARTMENT OF PHYSICS

Contact person:

Name: Jan Myrheim

Telephone: 73 59 36 53 (mobile 900 75 172)

Examination, course FY8304/FY3107
Mathematical approximation methods in physics

Wednesday December 3, 2014

Time: 09:00–13:00

Grades made public: Saturday January 3, 2015

Allowed to use: Calculator, mathematical tables.

All subproblems are given the same weight in the grading.

Problem 1:

Classification of a linear differential equation

Given the differential equation

$$y''(x) + \left(1 + \frac{x}{x^2 + 1}\right) y'(x) + \frac{1}{4} \left(1 + \frac{2x - 1}{x^2 + 1}\right) y(x) = 0 .$$

- a) Find the singular points of the equation, and classify them.
- b) Show that the leading asymptotic behaviour of $y(x)$ when $x \rightarrow \infty$ is:

$$y(x) \sim C_{\pm} x^{\pm \frac{1}{2}} e^{-\frac{x}{2}} ,$$

where C_{\pm} are constants.

- c) Find the leading asymptotic behaviour of $y(x)$ at the other singular points (two answers for every singular point).
- d) Can you find functions that have the correct asymptotic behaviour at all the singular points of the equation, and are analytical everywhere else?
Are they solutions of the equation?

Problem 2:**A boundary layer problem**

Given the boundary value problem

$$\epsilon y''(x) + e^{-x} y'(x) - e^{y(x)} = 0, \quad y(0) = 1, \quad y(1) = -1,$$

where ϵ is a small positive parameter. We calculate here to lowest order in ϵ .

- a) Where is the boundary layer?
How does the thickness of the boundary layer scale with ϵ ?
Give reasons for your answers.
- b) Find the outer solution (which is approximately valid outside the boundary layer).
- c) Find the inner solution (approximately valid inside the boundary layer).
- d) Find the uniform solution (approximately valid in the whole interval $[0, 1]$).
Sketch what it looks like.
- e) The WKB method can not be used for solving this boundary layer problem. Why not?

Problem 3:**Reduction of dimension**

We consider the wave equation in one time dimension and d space dimensions:

$$\left(\frac{\partial^2}{\partial t^2} - \nabla^2 \right) \phi(t, x_1, x_2, \dots, x_d) = 0,$$

where

$$\nabla^2 = \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \dots + \frac{\partial^2}{\partial x_d^2}.$$

Define

$$s = t^2 - x_1^2 - x_2^2 - \dots - x_d^2.$$

Assume that ϕ is a function of s alone, $\phi(t, x_1, x_2, \dots, x_d) = F(s)$.

- a) Derive an equation for $F(s)$, and solve it.
These Lorentz invariant solutions of the wave equation have no obvious physical interest.
- b) Show that the general solution of the wave equation in one space dimension is

$$\phi(t, x) = f(t - x) + g(t + x),$$

where f and g are arbitrary functions.

Is this consistent with your solution to problem 3a)?