


P1

2 sources, same band: 400-600 nm

$$\left. \begin{aligned} \lambda_1 &= 400 \text{ nm} = 0.4 \mu\text{m} \\ \lambda_2 &= 600 \text{ nm} = 0.6 \mu\text{m} \end{aligned} \right\}$$

SPECTRAL PHOTON
FLUX DENSITY:
 $\frac{\text{phot}}{\text{m}^2 \cdot \text{s} \cdot \mu\text{m}}$

$$\left\{ \begin{aligned} \uparrow_{\lambda A} &= \uparrow_A(\lambda) = A \lambda^3 \\ \uparrow_{\lambda B} &= \uparrow_B(\lambda) = B \lambda^{-2} \end{aligned} \right\}$$



Assume flat response: $\epsilon(\lambda) = \text{constant} (\equiv 1)$

Note:
 λ in microns, μm

Area of telescope: A_{TEL} (but not needed)

The two sources will "generate photoelectrons" at a rate ("count rate") R :

$$\frac{R_A}{A_{\text{TEL}}} = \int_{\lambda_1}^{\lambda_2} \uparrow_A(\lambda) d\lambda = A \int_{\lambda_1}^{\lambda_2} \lambda^3 d\lambda = A \left[\frac{\lambda^4}{4} \right]_{0.4}^{0.6} = \frac{0.6^4 - 0.4^4}{4} \cdot A \quad \left[\frac{\text{counts}}{\text{s} \cdot \text{cm}^2} \right]$$

(0.15) λ_1 (0.15) λ_2

$$\frac{R_B}{A_{\text{TEL}}} = \int_{\lambda_1}^{\lambda_2} \uparrow_B(\lambda) d\lambda = B \int_{\lambda_1}^{\lambda_2} \lambda^{-2} d\lambda = B \left[-\lambda^{-1} \right]_{0.4}^{0.6} = \left(\frac{1}{0.4} - \frac{1}{0.6} \right) B \quad \left[\frac{\text{counts}}{\text{s} \cdot \text{cm}^2} \right]$$

(0.15) λ_1 (0.15) λ_2

But they "generate photoelectrons at exactly the same rate":

$$(0.15) \quad R_A = R_B \Rightarrow \boxed{\frac{A}{B} = \frac{0.833}{0.026} = 32.05} \quad \left[\frac{1}{\mu\text{m}^5} \right]$$

("normalized" -0.17)

Brightness ratio F_B/F_A ? Now it's energy flux $\times h\nu = \frac{hc}{\lambda}$

$$F_A = \int_{\lambda_1}^{\lambda_2} \uparrow_A(\lambda) \cdot h\nu \cdot d\lambda = Ahc \int_{\lambda_1}^{\lambda_2} \lambda^2 d\lambda = Ahc \left[\frac{\lambda^3}{3} \right]_{0.4}^{0.6} = 0.05067 \cdot Ahc$$

(0.15)

$$F_B = \int_{\lambda_1}^{\lambda_2} \uparrow_B(\lambda) \cdot h\nu \cdot d\lambda = Bhc \int_{\lambda_1}^{\lambda_2} \lambda^{-3} d\lambda = Bhc \left[-\frac{1}{2\lambda^2} \right]_{0.4}^{0.6} = 1.736 \cdot Bhc$$

$$\Rightarrow \boxed{\frac{F_B}{F_A} = \frac{1.736 B}{0.05067 A} = 1.069}$$

(7% difference in F leads to the same R)
(wrong units but ok otherwise: 0.15)

NOTE:

(SPECTRAL) PHOTON FLUX
DENSITY

(SPECTRAL) ENERGY FLUX
DENSITY

$$f_\nu = \frac{\lambda^2}{c} f_\lambda$$

(15)

$$f(\lambda)$$

$$\times \frac{hc}{\lambda}$$

$$\rightarrow f_\lambda = f(\lambda) \text{ "f-lambda"}$$

$$f(\nu)$$


$$\times h\nu$$

$$\rightarrow f_\nu = f(\nu) \text{ "f-nu"}$$

IF you want to complicate things, this can be also done in ν space:

$$f(\lambda) \cdot h\nu \cdot \frac{\lambda^2}{c} = f_A(\nu) = \frac{A h c^4}{\nu^4}$$

$$; f_B(\nu) = \frac{B h}{c} \nu = f_B(\lambda) \cdot \lambda \nu \cdot \frac{\lambda^2}{c}$$




$$F_A = \int_{\nu_1}^{\nu_2} f_A(\nu) d\nu = \dots = \frac{A h c^4}{3} \cdot 0.15$$

$$F_B = \int_{\nu_1}^{\nu_2} f_B(\nu) d\nu = \dots = \frac{B h c}{2} \cdot 3.47$$

And reach the same $\frac{F_B}{F_A} = 1.069$

P2



DECLINATION: δ

ALTITUDE: h

ZENITH ANGLE: z

LATITUDE: L

a) 0,5p

$$\Rightarrow \boxed{h = 90 - L + \delta}$$

AT HIGHEST POINT:

$$\bullet h = 90 - L + \delta > 0^\circ$$

$$\Rightarrow \underline{\underline{\delta > L - 90^\circ = -40^\circ}}$$


$$\bullet h = 90 - L + \delta > 30^\circ$$

$$\Rightarrow \underline{\underline{\delta > L - 60^\circ = -10^\circ}}$$

b)

$$x \approx \frac{1}{\cos z} = 1,15$$

0,5p



K.EV ($L=50^\circ$)

above horizon: 30° :

$\delta > -40^\circ$ $\delta > -10^\circ$

$$\Rightarrow \theta = 90 + 10 = 100$$

$$(\theta < 100^\circ)$$

$$d\Omega = \sin\theta \, d\theta \, d\varphi$$

[if only, $\theta < 90 + 50 = 140^\circ$
 $\Rightarrow 0,57$]

VISIBLE SKY:
$$\Omega_{\text{vis}} = \int_{\text{vis}} d\Omega = \int_0^{2\pi} d\varphi \int_{\theta=0^\circ}^{\theta=\theta_{\text{max}}} \sin\theta \, d\theta = 2\pi (1 - \cos\theta_{\text{max}})$$

$$-\cos\theta \Big|_0^{\theta_{\text{max}}} = \cos 0 - \cos\theta_{\text{max}} = 1 - \cos\theta_{\text{max}}$$

FULL SKY: $\Omega_{\text{SKY}} = 4\pi$


\Rightarrow FRACTION:
$$\frac{\Omega_{\text{vis}}}{\Omega_{\text{SKY}}} = \frac{1 - \cos\theta_{\text{max}}}{2} = 0,587$$

58,7%

P3

$$d \circ \pi \left(\frac{d}{2}\right)^2$$


25 μm



$$\left. \begin{aligned} d &= 15'' \\ F &= 10^{-7} \text{ W m}^{-2} \end{aligned} \right\}$$

$$R = \frac{f_L}{D}$$

a) $B = \frac{F}{A} = \frac{4F}{\pi d^2} = 5,66 \times 10^{-10} \text{ W m}^{-2} \text{ arcsec}^{-2}$



b) $R = 8$ (f18) $f_L = D \cdot R = 0,3 \text{ m} \cdot 8 = 2,4 \text{ m}$ 0,25

AREA PIXEL : 25 μm \times 25 μm .

ANGULAR " " : $\Delta\theta \times \Delta\theta$ where $\Delta\theta = \frac{\Delta x}{f_L} = 1,0416 \times 10^{-5} \text{ rad}$

$$\times \frac{180^\circ}{\pi \text{ rad}} \times \frac{3600''}{1^\circ} \rightarrow \Delta\theta = 2,15''$$

$$E = B \times A_{\text{TEL}} \times A_{\text{PIX}} \times T = B \times \pi \left(\frac{D}{2}\right)^2 \times \Delta\theta^2 \times T = 1,85 \times 10^{-8} \text{ J}$$

↑
circular pixel - 0,25

$$c) \text{ New } \left. \begin{array}{l} D = 30 \text{ m} \\ R = 4 \end{array} \right\} \rightarrow \Delta f_L = D \cdot R = 120 \text{ m} \quad 0,25$$

$$\Delta \theta = \frac{\Delta x}{f_L} = 2,08 \times 10^{-7} \text{ rad} = 0,0429''$$

$$E = B \times A_{TEL} \times A_{PIX} \times T = 7,36 \times 10^{-8} \text{ J}$$

↑
window fixed -0,25

Q4


95,000 e⁻ < 100,000 e⁻ FULL-WELL DEPTH ✓

~~Pixel~~ PIXEL "COUNTS" = $\frac{95000 e^-}{1.4 e^-/ADU}$ = 67857 ADU

16-bit ADC: $2^{16} - 1 = \underline{65535 \text{ MAX}}$

COUNTS > MAX ⇒ SATURATED

Q7



b) X-rays penetrate matter (metals), reflected fraction drops drastically with increasing photon energy.

GRAZING INCIDENCE MIRRORS are used to focus X-rays, since reflection is still efficient at very small incident angles (<< 1°). This results in long focal lengths...

Q5

$$L_{\text{peak}} = (3,79 \pm 0,15) \times 10^{38} \text{ erg/s}$$

a)

[0,5]

$$F_{\text{peak}} = (1,27 \pm 0,13) \times 10^{-7} \text{ erg/cm}^2$$

$$F = \frac{L}{4\pi d^2} \Rightarrow d = \sqrt{\frac{L}{4\pi F}} = 1,54 \times 10^{22} \text{ cm} \\ = 5,0 \text{ kpc}$$

$$\delta d = \sqrt{\delta d_F^2 + \delta d_L^2}$$

$$\delta d_F = \frac{\partial d}{\partial F} \delta F = \frac{1}{2} \cdot \sqrt{\frac{L}{4\pi F}} \cdot \frac{\delta F}{F}$$

$$\delta d_L = \frac{\partial d}{\partial L} \delta L = \frac{1}{2} \sqrt{\frac{L}{4\pi F}} \frac{\delta L}{L}$$

$$\Rightarrow \frac{\delta d}{d} = \sqrt{\frac{1}{4} \left(\frac{\delta F}{F}\right)^2 + \frac{1}{4} \left(\frac{\delta L}{L}\right)^2} = \frac{1}{2} \sqrt{\left(\frac{\delta F}{F}\right)^2 + \left(\frac{\delta L}{L}\right)^2} = 0,0556 \\ \frac{1}{2} \sqrt{(0,104)^2 + (0,0396)^2} \quad (5,6\%)$$

$$\Rightarrow \delta d = 278 \text{ pc} = 0,278 \text{ kpc}$$

$$\Rightarrow d = 5,0 \pm 0,3 \text{ kpc} = 4,99 \pm 0,28 \text{ kpc}$$


0,25% 0,25%

b)
[0.5]

$$\frac{X - \bar{X}}{\sigma} = \frac{5,56 - 4,99}{0,28} = 2$$

"2 sigma":

Probab (> +2σ deviation) ∴



From Table 6.2: $P = \frac{0,0465}{2} = 0,0228$

~~2,8%~~

2,3%

factor 2 off: 0,25%

Correct reasoning but wrong σ: 0,5%

Q6:

$$\Delta = 30 \text{ m},$$

$$f/15 : R = \frac{f_L}{\Delta} = 15$$

a)
[0.5r]

$$f_L = 15 \times \Delta = 30 \times 15 = 450 \text{ m} = 450.000 \text{ mm}$$

$$\text{Plate scale: } \frac{1}{f_L} \rightarrow \frac{206265''}{f_L (\text{mm})} = 0,458 \frac{''}{\text{mm}}$$

b) [0.5r] $20 \times 60' = 1200'' \Rightarrow \text{FOV: } 1200'' \times 1200''$

$$\times \frac{1}{0,458 \frac{''}{\text{mm}}} \Rightarrow \text{FOV: } 2620 \text{ mm} \times 2620 \text{ mm}$$

$$\times \frac{1 \text{ pix}}{30 \times 10^{-3} \text{ mm}} \Rightarrow 87336 \times 87336 = \underline{\underline{7,63 \times 10^9 \text{ pix}}}$$

correct reasoning, wrong plate scale: 0,5r.

Q7

a) $R = 2850 = \frac{\lambda}{\Delta\lambda} \rightarrow \Delta\lambda = \frac{\lambda}{R} = 1,7 \text{ \AA}$


0,25

$\frac{\Delta\nu}{c} = \frac{\Delta\lambda}{\lambda} \rightarrow \Delta\nu = \frac{c}{R} = \frac{105,2}{105} \text{ km/s} \approx 1 \text{ km/s}$

vel. missing = -0,1

b) $\langle \text{disp} \rangle = \frac{5730 - 4000}{1018} = 1,7 \frac{\text{ \AA}}{\text{pix}}$

0,688 $\frac{\text{pix}}{\text{ \AA}}$ OK ✓



0,25

d) Nyquist: $V_{\text{sample}} \geq V_{\text{Nyq}}$ (≥ 2 pixels per resolution element)

0,25

⇒ (slightly) UNDER-SAMPLED