

Solutions to the final exam  
TFY4210 – Applied quantum mechanics  
May 23, 2007

**Problem 1**

- a) The functional  $F[n]$  is said to be *universal* because it is independent of the external potential  $v_{\text{ext}}$  and does not refer to a specific system. This implies that the same  $F[n]$  is to be used for *all* electron structure problems, whether the calculation involves only a single atom or a molecule adsorbed on a metal surface.
- b) Normalising:

$$\begin{aligned} N &= C \int d\mathbf{r} e^{-r/R} \\ &= 4\pi C \int_0^\infty dr r^2 e^{-r/R} \\ &= 4\pi C R^3 \int_0^\infty dt t^2 e^{-t} \\ &= 4\pi C R^3 \Gamma(3) \\ &= 8\pi C R^3 \end{aligned}$$

So  $C = N/(8\pi R^3)$ , and

$$n(r) = \frac{N}{8\pi R^3} e^{-r/R}$$

It is a good idea to check the units:  $[R] = \text{m}$ , so the electron (*number*) density has the correct unit  $[n] = \text{m}^{-3}$ .

- c) We first calculate  $E_{\text{ext}}$ :

$$\begin{aligned} E_{\text{ext}}(R) &= -\frac{Ze^2}{4\pi\epsilon_0} \int d\mathbf{r} \frac{n(r)}{r} \\ &= -\frac{NZe^2}{32\pi^2\epsilon_0 R^3} 4\pi \int_0^\infty dr r e^{-r/R} \\ &= -\frac{e^2}{8\pi\epsilon_0} \frac{NZ}{R^3} R^2 \int_0^\infty dt t e^{-t} \end{aligned}$$

$$\begin{aligned}
&= -\frac{e^2}{8\pi\epsilon_0} \frac{NZ}{R} \Gamma(2) \\
&= -\underbrace{\frac{e^2}{8\pi\epsilon_0}}_{\equiv C_1} \frac{NZ}{R}
\end{aligned}$$

(Note that  $[e^2/(\epsilon_0 R)] = \text{C}^2/(\text{C}^2 \text{J}^{-1} \text{m}^{-1} \text{m}) = \text{J}$ , so  $[E_{\text{ext}}] = \text{J}$ .)

Then the kinetic energy  $T_{\text{TF}}$ :

$$\begin{aligned}
T_{\text{TF}}(R) &= A_s \int d\mathbf{r} n^{5/3}(r) \\
&= A_s \frac{N^{5/3}}{(8\pi)^{5/3} R^5} 4\pi \int_0^\infty dr r^2 e^{-5r/3R} \\
&= A_s \frac{N^{5/3}}{(8\pi)^{5/3} R^5} 4\pi \left(\frac{3}{5}R\right)^3 \int_0^\infty dt t^2 e^{-t} \\
&= A_s \frac{N^{5/3}}{(8\pi)^{5/3} R^5} 4\pi \left(\frac{3}{5}R\right)^3 \Gamma(3) \\
&= \underbrace{\frac{27}{125} \frac{A_s}{(8\pi)^{2/3}}}_{\equiv C_2} \frac{N^{5/3}}{R^2}
\end{aligned}$$

(Units:  $A_s \propto \hbar^2/m_e$  so  $[A_s] = \text{J}^2 \text{s}^2/\text{kg} = \text{Jm}^2$  and  $[T_{\text{TF}}] = \text{Jm}^2 \text{m}^{-2} = \text{J}$ .)

We finally calculate the Hartree energy:

$$\begin{aligned}
E_{\text{H}}(R) &= \frac{e^2}{8\pi\epsilon_0} \int d\mathbf{r}_1 d\mathbf{r}_2 \frac{n(r_1)n(r_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} \\
&= \frac{e^2}{(8\pi)^3 \epsilon_0} \frac{N^2}{R^6} \int d\mathbf{r}_1 d\mathbf{r}_2 \frac{e^{-(r_1+r_2)/R}}{|\mathbf{r}_1 - \mathbf{r}_2|}
\end{aligned}$$

Introducing the new parameters  $\mathbf{s}_1 = R\mathbf{r}_1$  and  $\mathbf{s}_2 = R\mathbf{r}_2$ :

$$E_{\text{H}}(R) = \frac{e^2}{(8\pi)^3 \epsilon_0} \frac{N^2}{R} \int d\mathbf{s}_1 d\mathbf{s}_2 \frac{e^{-(s_1+s_2)}}{|\mathbf{s}_1 - \mathbf{s}_2|}$$

Then we use the expansion (given on page 5 of the exam set):

$$\frac{1}{|\mathbf{s}_1 - \mathbf{s}_2|} = \frac{1}{s_1} \sum_{l,m} \left(\frac{4\pi}{2l+1}\right) \left(\frac{s_2}{s_1}\right)^l Y_{lm}(\theta_1, \phi_1) Y_{lm}(\theta_2, \phi_2)$$

when  $s_2 > s_1$ , and with  $s_1$  and  $s_2$  interchanged when  $s_2 < s_1$ . Our trial electron density  $n(r)$  is independent of angles, so the integration over the angles is easy. Only the spherical harmonics depend on the angles, so

$$\int d\Omega_1 \int d\Omega_2 \frac{1}{|\mathbf{s}_1 - \mathbf{s}_2|} = \frac{1}{s_1} \sum_{l,m} \left( \frac{4\pi}{2l+1} \right) \left( \frac{s_2}{s_1} \right)^l \int d\Omega_1 Y_{lm}(\theta_1, \phi_1) \int d\Omega_2 Y_{lm}(\theta_2, \phi_2)$$

Since

$$\begin{aligned} \int d\Omega_1 Y_{lm}(\theta_1, \phi_1) &= \sqrt{4\pi} \int d\Omega_1 Y_{00}^*(\theta_1, \phi_1) Y_{lm}(\theta_1, \phi_1) \\ &= \sqrt{4\pi} \delta_{l,0} \delta_{m,0} \end{aligned}$$

we get

$$\int d\Omega_1 \int d\Omega_2 \frac{1}{|\mathbf{s}_1 - \mathbf{s}_2|} = \begin{cases} (4\pi)^2/s_1 & : s_2 > s_1 \\ (4\pi)^2/s_2 & : s_2 < s_1 \end{cases}$$

The radial integration must be split into two parts, one with  $s_2 < s_1$  and one with  $s_2 > s_1$ :

$$\begin{aligned} E_H(R) &= \frac{1}{(8\pi)^3 \varepsilon_0} \frac{e^2 N^2}{R} (4\pi)^2 \int_0^\infty ds_2 s_2^2 e^{-s_2} \left[ \int_0^{s_2} ds_1 \frac{s_1^2}{s_2} e^{-s_1} + \int_{s_2}^\infty ds_1 \frac{s_1^2}{s_1} e^{-s_1} \right] \\ &= \frac{1}{32\pi \varepsilon_0} \frac{e^2 N^2}{R} \int_0^\infty ds_2 s_2^2 e^{-s_2} \left[ \frac{2}{s_2} - \frac{1}{s_2} (2 + 2s_2 + s_2^2) e^{-s_2} + (1 + s_2) e^{-s_2} \right] \\ &= \frac{1}{32\pi \varepsilon_0} \frac{e^2 N^2}{R} \left[ 2 \int_0^\infty ds_2 s_2 e^{-s_2} - 2 \int_0^\infty ds_2 s_2 e^{-2s_2} - \int_0^\infty ds_2 s_2^2 e^{-2s_2} \right] \\ &= \frac{1}{32\pi \varepsilon_0} \frac{e^2 N^2}{R} \left[ 2\Gamma(2) - \frac{1}{2}\Gamma(2) - \frac{1}{8}\Gamma(3) \right] \\ &= \frac{5}{32} \left( \frac{e^2}{4\pi \varepsilon_0} \right) \frac{N^2}{R} \end{aligned}$$

$$(\text{Units: } [E_H] = [e^2/(\varepsilon_0 R)] = C^2/(C^2 J^{-1} m^{-1} m) = J)$$

d) We can find an approximation for the ground state energy by minimising

$$\begin{aligned} E(R) &= E_{\text{ext}}(R) + T_{\text{TF}}(R) + E_H(R) \\ &= -C_1 \frac{NZ}{R} + C_2 \frac{N^{5/3}}{R^2} + C_k \frac{N^2}{R} \\ &= -(C_1 Z - C_k N) \frac{N}{R} + C_2 \frac{N^{5/3}}{R^2} \end{aligned} \tag{1}$$

Here I have introduced

$$C_k = \frac{5}{32} \frac{1}{4\pi\epsilon_0}.$$

We first find the value  $R_{\min}$  that minimise  $E(R)$ :

$$\left. \frac{\partial E(R)}{\partial R} \right|_{R=R_{\min}} = 0 = (C_1 Z - C_k N) \frac{N}{R_{\min}^2} - 2C_2 \frac{N^{5/3}}{R_{\min}^3}$$

which gives us:

$$R_{\min} = \frac{2C_2 N^{2/3}}{C_1 Z - C_k N}$$

Substituting this value in equation (1), we get our approximation for the ground state energy as a function of  $N$  and  $Z$ :

$$\begin{aligned} E(N, Z) &= - (C_1 Z - C_k N) \frac{N}{R_{\min}} + C_2 \frac{N^{5/3}}{R_{\min}^2} \\ &= - \left[ (C_1 Z - C_k N) - C_2 N^{2/3} \frac{1}{R_{\min}} \right] \frac{N}{R_{\min}} \\ &= - \left[ (C_1 Z - C_k N) - \frac{1}{2} (C_1 Z - C_k N) \right] N^{1/3} \frac{(C_1 Z - C_k N)}{2C_2} \\ &= - \frac{1}{4} N^{1/3} \frac{(C_1 Z - C_k N)^2}{C_2} \end{aligned}$$

The approximation for the ground state electron density is:

$$\begin{aligned} n(r, N, Z) &= \frac{N}{8\pi R_{\min}^3} e^{-r/R_{\min}} \\ &= \frac{(C_1 Z - C_k N)^3}{64\pi N C_2^3} e^{-r(C_1 Z - C_k N)/(2C_2 N^{2/3})} \end{aligned}$$

e) Neutral atom,  $N = Z$ :

$$\begin{aligned} E_{\text{GS}}(Z, Z) &= - \frac{1}{4} Z^{1/3} \frac{(C_1 Z - C_k Z)^2}{C_2} \\ &= - \underbrace{\frac{1}{4} \frac{(C_1 - C_k)^2}{C_2}}_{\equiv C_3} Z^{7/3} \end{aligned}$$

Our value for  $C_3$  is

$$\begin{aligned}
C_3 &= \frac{1}{4} \frac{(C_1 - C_k)^2}{C_2} \\
&= \frac{1}{4} \frac{e^4}{16\pi^2 \varepsilon_0^2} \left( \frac{1}{2} - \frac{5}{32} \right)^2 \frac{125 (8\pi)^{2/3}}{27 A_s} \\
&= \frac{5^4 11^2}{2^{15} 3^4} \left( \frac{8}{3\pi} \right)^{2/3} \frac{e^4 m_e}{\pi^2 \varepsilon_0^2 \hbar^2} \\
&\approx 1.7818 \cdot 10^{-18} \text{ J}
\end{aligned}$$

Converted to the Rydberg unit:

$$C_3 = 0.81740 \text{ Ry}$$

So, with our choice of trial electron density we are almost 50% in error compared to the exact value. At least we have found an upper limit to the ground state energy (as expected for a variational approximation):

$$E_{\text{GS}}^{\text{approx}} = -0.81740 Z^{7/3} \text{ Ry}$$

g) For a neutral atom our approximation to the electron density is:

$$n(r, Z) = \frac{Z^2 \gamma^3}{8\pi} e^{-\gamma r Z^{1/3}}$$

where

$$\gamma = (C_1 - C_k)/(2C_2) = \frac{1375}{1728} \frac{e^2}{4\pi\varepsilon_0} \frac{(8\pi)^{2/3}}{A_s} \approx 4.4936 \cdot 10^{10} \text{ m}^{-1}$$

which is larger than zero, so our approximate electron density is an exponentially decreasing function of  $r$ . Figure 1 shows plots of the electron density (more precisely the radial distribution of electrons  $4\pi r^2 n(r)$ ) for Xe. The plot on the left hand side shows the result of a (much) more precise LDA (local density approximation) calculation (Note that they have used atomic units, so the values along the  $r$ -axis differ in the two plots) and on the right hand side is a plot of our electron density. As you can see, our electron density lacks the shell structure seen in the LDA-plot.

h) In our approximation we have

- *not* included the exchange energy-contribution to the total energy.




Figure 1:

- *not* included the electron correlation-contribution to the total energy.
- used a rather crude approximation for the kinetic energy (assumed that it is equal to the kinetic energy of a non-interacting electron gas (ideal Fermi gas)).

## Problem 2

a) Our starting point is

$$i\hbar \frac{\partial \Psi}{\partial t} = H \Psi$$

where  $H = c\sqrt{\mathbf{p}^2}$  for a massless particle. Can the square root be linearised?

$$\sqrt{\mathbf{p}^2} = \alpha_1 p_1 + \alpha_2 p_2 + \alpha_3 p_3$$

where  $\alpha_i$  (with  $i = 1, 2, 3$ ) are unknown quantities to be determined. Take the square of each side of this equation:

$$p_1^2 + p_2^2 + p_3^2 = \alpha_1^2 p_1^2 + \alpha_2^2 p_2^2 + \alpha_3^2 p_3^2 + \\ + (\alpha_1 \alpha_2 + \alpha_2 \alpha_1) p_1 p_2 + (\alpha_1 \alpha_3 + \alpha_3 \alpha_1) p_1 p_3 + (\alpha_3 \alpha_2 + \alpha_2 \alpha_3) p_2 p_3$$

The right and left hand side are equal if the quantities  $\alpha_i$  satisfy the *anti*-commutator relations

$$\alpha_i \alpha_j + \alpha_j \alpha_i \equiv \{\alpha_i, \alpha_j\} = 2\delta_{ij}$$

where  $\delta_{ij}$  is the Kroenecker delta. In addition  $\alpha_i$  must be hermitian  $\alpha_i^\dagger = \alpha_i$  (so that  $H$  is hermitian). Other properties:

1.  $\alpha_i^2 = 1$  so the eigen values of  $\alpha_i$  must be  $\pm 1$ .

2.  $\text{Tr } \alpha_i = 0$

Proof: (shown only for  $\alpha_1$ )

$$\text{Tr } \alpha_1 = \text{Tr}(\alpha_1 \alpha_2^2) = \text{Tr}(\alpha_2 \alpha_1 \alpha_2) = -\text{Tr}(\alpha_2^2 \alpha_1) = -\text{Tr } \alpha_1$$

which implies that  $\text{Tr } \alpha_1 = 0$ . Here I have used the fact that  $\alpha_1$  and  $\alpha_2$  anti-commute, and that the trace is cyclic (i.e.,  $\text{Tr}(ABC) = \text{Tr}(BCA) = \text{Tr}(CAB)$ , etc, for matrices  $A, B$  and  $C$ ).

3. The dimension of  $\alpha_i$  must be *even*, i.e.  $D = 2n$  where  $n = 1, 2, 3, \dots$ . This follows from the property  $\text{Tr } \alpha_i = 0$  and the fact that the eigen values to  $\alpha_i$  are  $\pm 1$  (the trace of a matrix is equal to the sum of its eigen values).

Thus, we get the following equation

$$i\hbar \frac{\partial \Psi}{\partial t} = H\Psi = c\boldsymbol{\alpha} \cdot \hat{\mathbf{p}}\Psi = \frac{c\hbar}{i}\boldsymbol{\alpha} \cdot \nabla\Psi$$

where  $\boldsymbol{\alpha} = (\alpha_1, \alpha_2, \alpha_3)$  are  $2n$ -dimensional matrices with the above mentioned properties. This is of course exactly the same equation as we get if we put  $m = 0$  in the Dirac equation.

b) We need *three* hermitian and mutually anti-commuting matrices. The Pauli matrices are three such matrices, so in this case we can choose  $\boldsymbol{\alpha} = \boldsymbol{\sigma}$ . We then get the following two equations (when we do not forget both signs  $E = \pm c\sqrt{\mathbf{p}}$ ):

$$\frac{\partial \Psi}{\partial t} = \pm c\boldsymbol{\sigma} \cdot \nabla\Psi$$

We *cannot* use the Pauli matrices when  $m \neq 0$ , because we then need *four* hermitian, anti-commuting matrices. Assume to the contrary that there is such a fourth matrix:

$$\tilde{\sigma} = \begin{pmatrix} a & b^* \\ b & -a \end{pmatrix}$$

This matrix is hermitian and  $\text{Tr } \tilde{\sigma} = 0$ . It must anti-commute with  $\sigma_z$ :

$$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} a & b^* \\ b & -a \end{pmatrix} = - \begin{pmatrix} a & b^* \\ b & -a \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

By multiplying the matrices we get

$$\begin{pmatrix} a & b^* \\ -b & a \end{pmatrix} = \begin{pmatrix} -a & b^* \\ -b & -a \end{pmatrix}$$

thus  $a = 0$ :

$$\tilde{\sigma} = \begin{pmatrix} 0 & b^* \\ b & 0 \end{pmatrix}$$

Our matrix must also anti-commute with  $\sigma_x$ . It is easy to show that this demand implies  $b^* = -b$ , so

$$\tilde{\sigma} = \text{constant} \times \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} = \text{constant} \times \sigma_y$$

and so  $\tilde{\sigma}$  will *not* anti-commute with  $\sigma_y$ . It is therefore not possible to find four mutually anti-commuting, hermitian  $2 \times 2$ -matrices, and we can therefore not use the Pauli matrices when  $m \neq 0$ .

c) Assume a plane-wave solution:

$$\Psi = u e^{-\frac{i}{\hbar}(Et - \mathbf{p} \cdot \mathbf{r})}$$

where  $u$  is a two-component spinor. Then we find from one of the Weyl equations:

$$Eu = c\boldsymbol{\sigma} \cdot \mathbf{p}u$$

Written on matrix form:

$$\begin{pmatrix} E + cp_z & c(p_x + ip_y) \\ c(p_x - ip_y) & E - cp_z \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = 0$$

This equation has non-trivial solutions only if the determinant of the system's matrix is zero:

$$(E + cp_z)(E - cp_z) - c^2(p_x + ip_y)(p_x - ip_y) = 0$$

which yields:

$$E^2 - c^2\mathbf{p}^2 = 0$$

or equivalently  $E = \pm pc$ .

### Problem 3

- a) –  $\mathbf{e}_{\mathbf{k},\lambda}$  are two ( $\lambda = 1, 2$ ) unit polarisation vectors normal to each other.  
 –  $a_{\mathbf{k},\lambda}^\dagger$  and  $a_{\mathbf{k},\lambda}$  are creation and annihilation operators.  $a_{\mathbf{k},\lambda}^\dagger$  creates a photon in the mode  $(\mathbf{k}, \lambda)$ , while  $a_{\mathbf{k},\lambda}$  destroys a photon in the mode  $(\mathbf{k}, \lambda)$ .

Coulomb gauge

$$0 = \nabla \cdot \mathbf{A} = i \sum_{\mathbf{k},\lambda} (\mathbf{k} \cdot \mathbf{e}_{\mathbf{k},\lambda}) \sqrt{\frac{\hbar}{2V\epsilon_0\omega_{\mathbf{k}}}} \left( a_{\mathbf{k},\lambda} e^{i\mathbf{k} \cdot \mathbf{r}} - a_{\mathbf{k},\lambda}^\dagger e^{-i\mathbf{k} \cdot \mathbf{r}} \right)$$



which implies

$$\mathbf{k} \cdot \mathbf{e}_{\mathbf{k},\lambda} = 0$$

so the polarisation vectors  $\mathbf{e}_{\mathbf{k},1}$  and  $\mathbf{e}_{\mathbf{k},2}$  are normal to the wave vector  $\mathbf{k}$ .

b) Now

$$\nabla \cdot (\mathbf{A}(\mathbf{r})f(\mathbf{r})) = (\nabla \cdot \mathbf{A}(\mathbf{r}))f(\mathbf{r}) + \mathbf{A}(\mathbf{r}) \cdot \nabla f(\mathbf{r})$$

so choosing Coulomb gauge ( $\nabla \cdot \mathbf{A} = 0$ ) we get

$$\nabla \cdot (\mathbf{A}(\mathbf{r})f(\mathbf{r})) = \mathbf{A}(\mathbf{r}) \cdot \nabla f(\mathbf{r})$$

thus  $\mathbf{p}$  and  $\mathbf{A}$  commute when we use the Coulomb gauge, so

$$\begin{aligned} V &= \frac{e}{2m} (\mathbf{A} \cdot \mathbf{p} + \mathbf{p} \cdot \mathbf{A}) + \frac{e^2}{2m} \mathbf{A}^2 \\ &= \frac{e}{2m} (2\mathbf{A} \cdot \mathbf{p}) + \frac{e^2}{2m} \mathbf{A}^2 \\ &= \frac{e}{m} \mathbf{A} \cdot \mathbf{p} + \frac{e^2}{2m} \mathbf{A}^2 \end{aligned}$$

c) In the case of spontaneous emission there is one extra photon in the final state compared to the initial state. To *lowest* order we therefore must include the term which is linear in the creation operator  $a_{\mathbf{k},\lambda}^\dagger$ , i.e.  $e/m\mathbf{A} \cdot \mathbf{p}$ :

$$\begin{aligned} M &\propto \langle \psi_f | \langle \dots, n_{\mathbf{k},\lambda} + 1, \dots | \frac{e}{m} \mathbf{A} \cdot \mathbf{p} | \dots, n_{\mathbf{k},\lambda}, \dots \rangle | \psi_i \rangle \\ &\propto \underbrace{\langle \dots, n_{\mathbf{k},\lambda} + 1, \dots | a_{\mathbf{k},\lambda}^\dagger | \dots, n_{\mathbf{k},\lambda}, \dots \rangle}_{\sqrt{n_{\mathbf{k},\lambda} + 1} \langle \dots, n_{\mathbf{k},\lambda} + 1, \dots | \dots, n_{\mathbf{k},\lambda} + 1, \dots \rangle = \sqrt{n_{\mathbf{k},\lambda} + 1}} \end{aligned}$$

d) In the electric dipole approximation we have the selection rule  $\Delta l = \pm 1$ , thus the transitions

$$\begin{aligned} 4p &\rightarrow 3d \\ &\rightarrow 3s \\ &\rightarrow 2s \\ &\rightarrow 1s \end{aligned}$$

are allowed.