

Formalities.


Solutions should be emailed latest Friday 27.03, at 15.00.

Executive summary.


Write a 3–4 page executive summary of the lectures (ending [and including] with spin 1/2). Omit derivations.

Higgs decay into fermions and the optical theorem.

In the Standard Model, the Higgs particle h is a scalar particle that interacts with all fermions via a Yukawa coupling y proportional to the fermion mass m , $y = \frac{1}{2} gm/m_W$,



- a.) Calculate the decay width $\Gamma(h \rightarrow \bar{f}f)$ of a Higgs particle with mass M into a antifermion-fermion pair (at tree-level).
- b.) Show that a fermion loop leads to an additional minus sign in the Feynman amplitude.
- c.) Consider the following contribution of fermions to the self-energy $\Sigma(p^2)$ of the Higgs,



Use dimensional regularisation to calculate $\Sigma^{\bar{f}f}$ and show that

$$\Sigma^{\bar{f}f} = \frac{A}{\epsilon} + B \left[C + \int_0^1 dz a^2 \ln(a^2/\mu^2) \right]$$

with $a^2 = m^2 - p^2 z(1 - z) - i\epsilon$. Note: In d spacetime dimensions, the Clifford algebra becomes $\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu} I_d$ with I_d as the d -dimensional unit matrix. Thus contractions change to

$$\gamma^\mu \gamma_\mu = dI_d, \quad \gamma^\mu \not{a} \gamma_\mu = (2 - d)\not{a}, \quad \gamma^\mu \not{a} \not{b} \gamma_\mu = 4a \cdot b I_d - (d - 4)\not{a} \not{b}. \quad (1)$$

However, it is standard to define $\text{tr}(I_d) = 4$, which has the advantage that trace relations like $\text{tr}[\not{a} \not{b}] = 4a \cdot b$ are unchanged.

- d.) Determine the imaginary part $\Im \Sigma^{\bar{f}f}$ of the self-energy and show that the optical theorem holds, i.e. that $\Im \Sigma^{\bar{f}f} = M\Gamma(h \rightarrow \bar{f}f)$ for $p^2 = M^2$.
- e.) Obtain $\Im \Sigma^{\bar{f}f}$ directly by “cutting the self-energy”: Consider

$$i\Sigma^{\bar{f}f}(p^2) = \int \frac{d^4q}{(2\pi)^4} \dots$$

for $p = (M, \mathbf{0})$; find the poles and apply the identity

$$\frac{1}{x \pm i\varepsilon} = P\left(\frac{1}{x}\right) \mp i\pi\delta(x)$$

to the q^0 integral in order to obtain the imaginary part.

Good luck!